

Phospho-AMPK α 1-S485 Rabbit pAb

货号: **AYP11929**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Rattus norvegicus , Homo sapiens , Mus musculus IHC: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	64kDa/65kDa
实测分子量	56kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic phosphorylated peptide around S485 of human PRKAA1 (NP_006242.5).
序列	SGSVS

靶点信息

研究背景	The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio . AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protect s cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alt ernatively spliced transcript variants encoding distinct isoforms have been observed.
基因ID	5562

基因名	PRKAA1
Swiss	Q13131
别名	PRKAA1;AMPK;AMPK α 1

产品验证



Western blot analysis of Phospho-AMPK α 1-S485 expressed in 293T using Phospho-AMPK α 1-S485 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time : 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn