

SFRP2 Rabbit pAb

货号: **AYP11940**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Crataegus , Homo sapiens
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
理论分子量	33kDa
实测分子量	35KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse heart
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 25-295 of human SFRP2 (NP_003004.1).
序列	LFLFGQPDFS YKRSNCKPI PANLQLCHGIEYQNMRLPNLLGHETMKEVLEQAGAWIPLVMKQCHPDTKKFLCSLFAPVCLD DLDETIQPCHSLCVQVKDRCAPVMSAFGFPWPDMLECDRFPQDNDLCIPLASSDHLLPATEEAPKVCEACKNKND DDDND IMETLCKNDFALKIKVKEITYINRDTKIILETKSKTIYKLNQVSRDLKKSVLWLKDSLQCTCEEMNDINAPYLVMGQKQG GEL VITSVKRWQKQGREFKRISRSIRKLQC

靶点信息

研究背景	This gene encodes a member of the SFRP family that contains a cysteine-rich domain homologous to the putative Wnt-binding site of Frizzled proteins. SFRPs act as soluble modulators of Wnt signaling. Methylation of this gene is a potential marker for the presence of colorectal cancer.
------	--

基因ID	6423
基因名	SFRP2
Swiss	Q96HF1
别名	SFRP2;FRP-2;SARP1;SDF-5

产品验证



Western blot analysis of SFRP2 expressed in Mouse kidney, Mouse heart using SFRP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn