

# VEGFC Rabbit pAb

货号: **AYP11956**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>IHC:</b> Homo sapiens <b>WB:</b> Homo sapiens , Oryctolagus cuniculus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	46kDa
实测分子量	47KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,K-562(Negative control)
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200-300 of human VEGFC (NP_005420.1).
序列	TISFANHTSCRCMSKLDVYRQVHSIIRSLPATLPQCQAANKTCPTNYMWNHICRCLAQEDFMFSSDAGDDSTDGFHDI CGPNKELDEETCQCVCRAGLR

## 靶点信息

研究背景	The protein encoded by this gene is a member of the platelet-derived growth factor/vascular endothelial growth factor (PDGF/VEGF) family. The encoded protein promotes angiogenesis and endothelial cell growth, and can also affect the permeability of blood vessels. The proprotein is further cleaved into a fully processed form that can bind and activate VEGFR-2 and VEGFR-3 receptors.
基因ID	7424

基因名	VEGFC
Swiss	P49767
别名	VEGFC;Fit4-L;LMPH1D;VRP

## 产品验证



Western blot analysis of VEGFC expressed in U-251MG,K-562(Negative control) using VEGFC Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)