

MYRF Rabbit pAb

货号: **AYP11962**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus IHC: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	120kDa/124kDa
实测分子量	124kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2
细胞定位	Cytoplasm,Endoplasmic reticulum membrane,Nucleus,Single-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human MYRF (NP_001120864.1).
序列	MEVVDTEALQRFFEGHDINGALEPSNIDTSILEEYISKEDASDLCFDPDISAPASSASYSHGQPAMPGSSGVHHLSPGGG PSPGRHGPLPPPGYGTPLN

靶点信息

研究背景	This gene encodes a transcription factor that is required for central nervous system myelination and may regulate oligodendrocyte differentiation. It is thought to act by increasing the expression of genes that effect myelin production but may also directly promote myelin gene expression. Loss of a similar gene in mouse models results in severe demyelination. Alternative splicing results in multiple transcript variants.
基因ID	745

基因名	MYRF
Swiss	Q9Y2G1
别名	MYRF;C11orf9;MRF;Ndt80;pqn-47

产品验证



Western blot analysis of MYRF expressed in HeLa,HepG2 using MYRF Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn