

# MT-ND2 Rabbit pAb

货号: **AYP12090**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens , Mus musculus, Homo sapiens
应用	<a href="#">WB</a> <a href="#">IHC</a> <a href="#">IF/ICC</a>
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	43kDa
实测分子量	39KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Rat brain
细胞定位	mitochondrial inner membrane,mitochondrial respiratory chain complex I
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 248-347 of human MT-ND2 (YP_003024027.1).
序列	LLSLGGLPPLTGFLPKWAIIEEFTKNNSLIPTIMATITLLNLYFYLRLLIYSTSITLLPMSNNVKMKWQFEHTKPTPFLPTLIALTTL LLPISPFMLMIL

## 靶点信息

研究背景	Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor. Essential for the catalytic activity and assembly of complex I.
基因ID	4536

基因名	MT-ND2
Swiss	P03891
别名	MTND2;MT-ND2

## 产品验证



Western blot analysis of MT-ND2 expressed in Mouse brain,Rat brain using MT-ND2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)