

— ABLYBIO, Help Your Research



PCK1 Rabbit pAb

货号: **AYP12158**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Primary Mouse liver , HEK293T,MCF7 , Mus musculus
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200 IP: 1:500 - 1:1000
理论分子量	34kDa/69kDa
实测分子量	69KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Mouse liver,Mouse kidney,Rat liver,Rat pancreas,Rat kidney
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 490-622 of human PCK1 (NP_002582.3).
------	--

靶点信息

研究背景	This gene is a main control point for the regulation of gluconeogenesis. The cytosolic enzyme encoded by this gene, along with GTP, catalyzes the formation of phosphoenolpyruvate from oxaloacetate, with the release of carbon dioxide and GDP. The expression of this gene can be regulated by insulin, glucocorticoids, glucagon, cAMP, and diet. Defects in this gene are a cause of cytosolic phosphoenolpyruvate carboxykinase deficiency. A mitochondrial isozyme of the encoded protein also has been characterized.
基因ID	5105
基因名	PCK1
Swiss	P35558 (https://www.uniprot.org/uniprotkb/P35558/entry)
别名	PCK1,PEPCK-C,PEPCK1,PEPCKC,PCK1 Rabbit pAb,Serine-protein kinase PCK1

产品验证



Western blot analysis of PCK1 expressed in Mouse liver,Rat liver,HepG2 using PCK1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)