

— ABLYBIO, Help Your Research



NUR77 Rabbit pAb

货号: **AYP12243**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	33kDa/64kDa/65kDa
实测分子量	64-80KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,SK-BR-3,U-937(negative)
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 300-400 of human NUR77 (NP_002126.2).
------	--

靶点信息

研究背景	This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. Expression is induced by phytohemagglutinin in human lymphocytes and by serum stimulation of arrested fibroblasts. The encoded protein acts as a nuclear transcription factor. Translocation of the protein from the nucleus to mitochondria induces apoptosis. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	3164
基因名	NR4A1
Swiss	P22736 (https://www.uniprot.org/uniprotkb/P22736/entry)
别名	NR4A1,GFRP1,HMR,N10,NAK-1,NGFIB,NP10,NUR77,TR3,NUR77 Rabbit pAb,Early response protein NAK1,Nuclear hormone receptor NUR/77,Orphan nuclear receptor HMR,Orphan nuclear receptor TR3,ST-59,Testicular receptor 3,NAK1

产品验证



Western blot analysis of NUR77 expressed in MCF7, SK-BR-3, U-937 (negative) using NUR77 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)