

ADCY2 Rabbit pAb

货号: **AYP12267**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	103kDa/123kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,Rat eye
细胞定位	Cell membrane,Cytoplasm,Membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 430-600 of human ADCY2 (NP_065433.2).
序列	VTLEHLNGAYKVEEGDGDIRDYPYLKQHLVKTYFVINPKGERRSPQHLFRPRHTLDGAKMRASVRMTRYLESWGAAKPFALHHRDMMTTENGKISTTDVPMGQHNFNQNRTRLRTKSQKKRFEELNERMIQAIDGINAQKQWLKSEDIQRISLLFYNKVLEKEYRATALPAF

靶点信息

研究背景	This gene encodes a member of the family of adenylate cyclases, which are membrane-associated enzymes that catalyze the formation of the secondary messenger cyclic adenosine monophosphate (cAMP). This enzyme is insensitive to Ca(2+)/calmodulin, and is stimulated by the G protein beta and gamma subunit complex.
基因ID	108

基因名	ADCY2
Swiss	Q08462
别名	ADCY2;AC2;HBAC2

产品验证



Western blot analysis of ADCY2 expressed in SH-SY5Y, Rat eye using ADCY2 Rabbit pAb at 1:1000. Second ary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn