

— ABLYBIO, Help Your Research



PGAM2 Rabbit pAb

货号: **AYP12290**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	28kDa
实测分子量	29kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	22Rv1,SH-SY5Y,U-87MG,Mouse kidney,Mouse testis,Mouse skeletal muscle,Rat brain,Rat kidney,Rat skeletal muscle
细胞定位	cytosol,extracellular exosome,nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-253 of human PGAM2 (NP_000281.2).
------	---

靶点信息

研究背景	Phosphoglycerate mutase (PGAM) catalyzes the reversible reaction of 3-phosphoglycerate (3-PGA) to 2-phosphoglycerate (2-PGA) in the glycolytic pathway. The PGAM is a dimeric enzyme containing, in different tissues, different proportions of a slow-migrating muscle (MM) isozyme, a fast-migrating brain (BB) isozyme, and a hybrid form (MB). This gene encodes muscle-specific PGAM subunit. Mutations in this gene cause muscle phosphoglycerate mutase deficiency, also known as glycogen storage disease X.
基因ID	5224
基因名	PGAM2
Swiss	P15259 (https://www.uniprot.org/uniprotkb/P15259/entry)
别名	PGAM2,GSD10,PGAM-M,PGAMM,PGAM2 Rabbit pAb,BPG-dependent PGAM 2,Muscle-specific phosphoglycerate mutase,Phosphoglycerate mutase isozyme M

产品验证



Western blot analysis of PGAM2 expressed in Mouse kidney, Rat brain, 22Rv1 using PGAM2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)