

# YTHDC1 Rabbit pAb

货号: **AYP12310**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens
应用	<a href="#">WB</a> <a href="#">IF/ICC</a>
推荐浓度	<b>WB:</b> 1:1000 - 1:5000 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	82kDa/84kDa
实测分子量	110KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HCT116,293F,RAW264.7,C6,Rat brain
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-74 of human YTHDC1 (NP_001026902.1).
序列	MAADSREEKDGELNVLDDILTEVPEQDDELYNPESEQDKNEKKGSKRKRSDRMESTDTKRQKPSVHSRQLVSKPL

## 靶点信息

研究背景	Regulator of alternative splicing that specifically recognizes and binds N6-methyladenosine (m6A-containing RNAs. M6A is a modification present at internal sites of mRNAs and some non-coding RNAs and plays a role in the efficiency of mRNA splicing, processing and stability. Acts as a key regulator of exon-inclusion or exon-skipping during alternative splicing via interaction with mRNA splicing factors SRSF3 and SRSF10. Specifically binds m6A-containing mRNAs and promotes recruitment of SRSF3 to its mRNA-binding elements adjacent to m6A sites, leading to exon-inclusion during alternative splicing. In contrast, interaction with SRSF3 prevents interaction with SRSF10, a splicing factor that promotes exon skipping: this prevents SRSF10 from binding to its mRNA-binding sites close to m6A-containing regions, leading to inhibit exon skipping during alternative splicing.
基因ID	91746
基因名	YTHDC1
Swiss	Q96MU7
别名	YTHDC1;YT521;YT521-B

## 产品验证



Western blot analysis of YTHDC1 expressed in Rat brain, HeLa, HCT116 using YTHDC1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)