

ADSL Rabbit pAb

货号: AYP12345

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Rattus norvegicus , Homo sapiens
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:20 - 1:100 IP: 1:50 - 1:100
理论分子量	48kDa/54kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,MCF7
细胞定位	cytosol
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-310 of human ADSL (NP_000017.1).
序列	MAAGGDHGSPDSYRSPLASRYASPEMCFVFSDDRYKFRTRWRQLWLWLAEAEQTLGLPITDEQIQEMKSNLENIDFKMAAE EEKRLRHDMVMAHVHTFGHCCPKAAGIIHLGATSCYVGDNTDLILRNALDLLLLPKLARVISRLADFAKERASLPTLGFTHFQP AQLTTVGKRCCLWIQDLCMDLQNLKRVRRDDLRFKRVKGTQASFLQLFEGDDHKVEQLDKMVTEKAGFKRAFIITGQ TYTRKVDIEVLSVLASLGASVHKICTDIRLLANLKEMEEPFEKQIQSSAMPYKRNPMSRERCCSLARH

靶点信息

研究背景	The protein encoded by this gene belongs to the lyase 1 family. It is an essential enzyme involved in purine metabolism, and catalyzes two non-sequential reactions in the de novo purine biosynthetic pathway: the conversion of succinylaminoimidazole carboxamide ribotide (SAICAR) to aminoimidazole carboxamide ribotide (AICAR) and the conversion of adenylosuccinate (S-AMP) to adenosine monophosphate (AMP). Mutations in this gene are associated with adenylosuccinase deficiency (ADSLD), a disorder marked with psychomotor retardation, epilepsy or autistic features. Alternatively spliced transcript variants have been found for this gene.
基因ID	158
基因名	ADSL
Swiss	P30566
别名	ADSL;AMPS;ASASE;ASL

产品验证



Western blot analysis of ADSL expressed in HepG2, MCF7 using ADSL Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn