

EHHADH Rabbit pAb

货号: **AYP12371**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Gallus gallus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	69kDa/79kDa
实测分子量	79kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,A-549,HeLa,HepG2,Mouse small intestine,Rat liver,Rat kidney
细胞定位	Peroxisome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 444-723 of human EHHADH (NP_001957.2).
序列	EVIPSQYSSPTTIATVMNLSKKIKKIGVVVGNCFGVGNRMLNPYYNQAYFLLEEGSKPEEVDQVLEEFGFKMGPFVSDLA GLDVGWKSRRKGQGLTGPTLLPGTPARKRGNRRYCPIDVLCELGRFGQKTGKGWYQYDKPLGRIHKPDPWLSKFLSRYR KTHHIEPRTISQDEILERCLYSLINEAFRILGEGIAASPEHIDVVYLHGYGWPRHKGGPMPFYASTVGLPTVLEKLQKYRQNP IPQLEPSDYLLKSLASQGNPPLKEWQSLAGSPSSKL

靶点信息

研究背景	The protein encoded by this gene is a bifunctional enzyme and is one of the four enzymes of the peroxisomal beta-oxidation pathway. The N-terminal region of the encoded protein contains enoyl-CoA hydratase activity while the C-terminal region contains 3-hydroxyacyl-CoA dehydrogenase activity. Defects in this gene are a cause of peroxisomal disorders such as Zellweger syndrome. Two transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	1962
基因名	EHHADH
Swiss	Q08426
别名	EHHADH;ECHD;FRTS3;L-PBE;LBFP;LBP;PBFE

产品验证



Western blot analysis of EHHADH expressed in Mouse small intestine, Rat liver, SW480 using EHHADH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn