

ApoM Rabbit pAb

货号: **AYP12386**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	13kDa/21kDa
实测分子量	21KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Human serum
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 22-188 of human ApoM (NP_061974.2).
序列	QCPEHSQLTTLGVDGKEFPEVHLGQWYFIAGAAPTKEELATFDVPDNI VFNMAAGSAPMQLHLRATIRMKDGLCVPRKWI YHLTEGSTDLRTEGRPDMKTELFSSSSCPGGIMLNETGGYQRFLLYNRSPHPPEKCVVEEFKSLTSCLD SKAFLLTPRNQEA CELSNN

靶点信息

研究背景	The protein encoded by this gene is an apolipoprotein and member of the lipocalin protein family. It is found associated with high density lipoproteins and to a lesser extent with low density lipoproteins and triglyceride-rich lipoproteins. The encoded protein is secreted through the plasma membrane but remains membrane-bound, where it is involved in lipid transport. Alternate splicing results in both coding and non-coding variants of this gene.
基因ID	55937
基因名	APOM
Swiss	O95445
别名	APOM;G3a;HSPC336;NG20;apo-M

产品验证



Western blot analysis of ApoM expressed in HepG2, Human serum using ApoM Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn