

# HAS1 Rabbit pAb

货号: **AYP12400**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus , Homo sapiens
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	64kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,DU145,Mouse skeletal muscle
细胞定位	Membrane,Multi-pass membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 80-390 of human HAS1 (NP_001514.2).
序列	RVAAAARGPLDAATARSVALTISAYQEDPAYLRQCLASARALLYPRARLRVLMVVDGNRAEDLYMVDMFREVFADEDPAT YVWDGNYHQPWEPAAAAGAVGAGAYREVEAEDPGRlavealvrtrrcvcvaqRWGGKREVMYAFKALGDSVDYVQVC DSDTRLDPMALLELVRVLDEDPRVGAVGGDVRILNPLDSWVSFLSSLRYWVAFNVERACQSYFHCVSCISGPLGLYRNNL LQQFLEAWYNQKFLGTHCTFGDDRHLTNRMLSMGYATKYTSRRCYSETPSSFLRWLSQQTRWSKSYFREWLYN

## 靶点信息

研究背景	Hyaluronan or hyaluronic acid (HA) is a high molecular weight unbranched polysaccharide synthesized by a wide variety of organisms from bacteria to mammals, and is a constituent of the extracellular matrix. HA is synthesized by membrane-bound synthase at the inner surface of the plasma membrane, and the chains are extruded through pore-like structures into the extracellular space. It serves a variety of functions, including space filling, lubrication of joints, and provision of a matrix through which cells can migrate. HA is actively produced during wound healing and tissue repair to provide a framework for ingrowth of blood vessels and fibroblasts. Changes in the serum concentration of HA are associated with inflammatory and degenerative arthropathies such as rheumatoid arthritis. In addition, the interaction of HA with the leukocyte receptor CD44 is important in tissue-specific homing by leukocytes, and overexpression of HA receptors has been correlated with tumor metastasis. HAS1 is a member of the newly identified vertebrate gene family encoding putative hyaluronan synthases, and its amino acid sequence shows significant homology to the hasA gene product of <i>Streptococcus pyogenes</i> , a glycosaminoglycan synthetase (DG42) from <i>Xenopus laevis</i> , and a recently described murine hyaluronan synthase. Alternative splicing results in multiple transcript variants.
基因ID	3036
基因名	HAS1
Swiss	Q92839
别名	HAS1;HAS

## 产品验证



Western blot analysis of HAS1 expressed in U-87MG, DU145, Mouse skeletal muscle using HAS1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)