

Cullin 4A Rabbit pAb

货号: **AYP12421**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Mus musculus Co-IP: Homo sapiens,Mus musculus
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100 IP: 1:500 - 1:1000
理论分子量	76kDa/87kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,BT-474,HeLa,Mouse testis
细胞定位	nucleoplasm,nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200-300 of human Cullin 4A (Q13619).
序列	IERERSGEAVDRSLLRSLGMLSDLQVYKDSFELKFLLEETNCLYAAEQRLMQEREVPEYLNHVSKRLEEEGDRVITYLDHS TQKPLIACVEKQLLGEHLT

靶点信息

研究背景	CUL4A is the ubiquitin ligase component of a multimeric complex involved in the degradation of DNA damage-response proteins (Liu et al., 2009 [PubMed 19481525]).[supplied by OMIM, Oct 2009]
基因ID	8451

基因名	CUL4A
Swiss	Q13619
别名	CUL4A;cullin-4A

产品验证



Western blot analysis of Cullin 4A expressed in Mouse testis, K-562, BT-274 using Cullin 4A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn