

— ABLYBIO, Help Your Research



BAG3 Rabbit pAb

货号: **AYP12535**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: Rattus norvegicus WB: Rattus norvegicus IHC: Rattus norvegicus Co-IP: Rattus norvegicus
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100 IP: 1:50 - 1:100
理论分子量	61kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,HeLa,HepG2,Mouse brain,Mouse testis
细胞定位	cytoplasm,cytosol,nucleus,Z disc
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 380-575 of human BAG3 (NP_004272.2).
------	--

靶点信息

研究背景	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner.
基因ID	9531
基因名	BAG3
Swiss	O95817 (https://www.uniprot.org/uniprotkb/O95817/entry)
别名	BAG3,BAG-3,BIS,CAIR-1,MFM6,BAG3 Rabbit pAb,Bcl-2-associated athanogene 3,Bcl-2-binding protein Bis, Docking protein CAIR-1

产品验证



Western blot analysis of BAG3 expressed in Mouse brain,A-549,HeLa using BAG3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)