

— ABLYBIO, Help Your Research



DLGAP5 Rabbit pAb

货号: **AYP12565**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Homo sapiens IF: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	85kDa/94kDa/95kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF-7,DU-145,K-562
细胞定位	Cytoplasm,Nucleus,cytoskeleton,spindle
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 547-846 of human DLGAP5 (NP_055565.3).
------	--

靶点信息

研究背景	Potential cell cycle regulator that may play a role in carcinogenesis of cancer cells. Mitotic phosphoprotein regulated by the ubiquitin-proteasome pathway. Key regulator of adherens junction integrity and differentiation that may be involved in CDH1-mediated adhesion and signaling in epithelial cells.
基因ID	9787
基因名	DLGAP5
Swiss	Q15398 (https://www.uniprot.org/uniprotkb/Q15398/entry)
别名	DLGAP5,DLG7,HURP,DLGAP5 Rabbit pAb,Discs large homolog 7,Disks large-associated protein DLG7,Hepato ma up-regulated protein,KIAA0008

产品验证



Western blot analysis of DLGAP5 expressed in MCF-7,DU-145,K-562 using DLGAP5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)