

— ABLYBIO, Help Your Research



# HHLA2 Rabbit pAb

货号: **AYP12580**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>IHC:</b> Homo sapiens , Mus musculus <b>WB:</b> Mus musculus
应用	WB
推荐浓度	<b>WB:</b> 1:1000 - 1:5000 <b>IHC:</b> 1:50 - 1:200
理论分子量	39kDa/46kDa
实测分子量	70-80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	A-204, HepG2 (Negative control), MDA-MB-231
细胞定位	Membrane, Single-pass type I membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-200 of human HHLA 2 (NP_009003.1).
------	---

## 靶点信息

研究背景	This gene encodes a protein ligand found on the surface of monocytes. The encoded protein is thought to regulate cell-mediated immunity by binding to a receptor on T lymphocytes and inhibiting the proliferation of these cells. Alternate splicing results in multiple transcript variants.
基因ID	11148
基因名	HHLA2
Swiss	Q9UM44 ( <a href="https://www.uniprot.org/uniprotkb/Q9UM44/entry">https://www.uniprot.org/uniprotkb/Q9UM44/entry</a> )
别名	HHLA2,B7-H5,B7-H7,B7H7,B7y,HHLA2 Rabbit pAb,Human endogenous retrovirus-H long terminal repeat-associated protein 2

## 产品验证



Western blot analysis of HHLA2 expressed in A-204, HepG2 (Negative control), MDA-MB-231 using HHLA2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))