

HHLA2 Rabbit pAb

货号: **AYP12580**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Homo sapiens , Mus musculus WB: Mus musculus
应用	WB
推荐浓度	WB: 1:1000 - 1:5000 IHC: 1:50 - 1:200
理论分子量	39kDa/46kDa
实测分子量	70-80KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-204,HepG2(Negative control),MDA-MB-231
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-200 of human HHLA2 (NP_009003.1).
序列	IYVPMNEQIVIGRLDEDIILPSSFERGGSEVVIHWKYQDSYKVVHSYYKGSDDLHLESQDPRYANRTSLFYNEIQNGNASLFFRRV SLLDEGIYTCYVGTAIQVITNKVVLKVGVFLTPVMKYEKRNTNSFLICSVLSVYPRPIITWKMDNTPISENNMEETGSLDSFSI NSPLN

靶点信息

研究背景	This gene encodes a protein ligand found on the surface of monocytes. The encoded protein is thought to regulate cell-mediated immunity by binding to a receptor on T lymphocytes and inhibiting the proliferation of these cells. Alternate splicing results in multiple transcript variants.
------	--

基因ID	11148
基因名	HHLA2
Swiss	Q9UM44
别名	HHLA2;B7-H5;B7-H7;B7H7;B7y

产品验证



Western blot analysis of HHLA2 expressed in A-204, HepG2 (Negative control), MDA-MB-231 using HHLA2 R abbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 3 0ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn