

— ABLYBIO, Help Your Research



# SIRT2 Rabbit pAb

货号: **AYP12604**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens,Rattus norvegicus <b>IHC:</b> Rattus norvegicus
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	30kDa/35kDa/39kDa/41kDa/43kDa
实测分子量	39KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	Cell projection,Chromosome,Cytoplasm,Midbody,Myelin membrane,Nucleus,Perikaryon,centriole,centrosome,cytoskeleton,growth cone,microtubule organizing center,perinuclear region,spindle
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 300-389 of human SIRT2 (NP_036369.2).
------	--

## 靶点信息

研究背景	This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Several transcript variants are resulted from alternative splicing of this gene.
基因ID	22933
基因名	SIRT2
Swiss	Q8IXJ6 ( <a href="https://www.uniprot.org/uniprotkb/Q8IXJ6/entry">https://www.uniprot.org/uniprotkb/Q8IXJ6/entry</a> )
别名	SIRT2,SIR2,SIR2L,SIR2L2,sirtuin 2,SIRT2 Rabbit pAb,NAD-dependent protein deacylase sirtuin-2,NAD-dependent protein defatty-acylase sirtuin-2,Regulatory protein SIR2 homolog 2,SIR2-like protein 2

## 产品验证



Western blot analysis of SIRT2 expressed in Rat brain using SIRT2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)