

SIRT2 Rabbit pAb

货号: **AYP12604**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens,Rattus norvegicus IHC: Rattus norvegicus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	30kDa/35kDa/39kDa/41kDa/43kDa
实测分子量	39KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	Cell projection,Chromosome,Cytoplasm,Midbody,Myelin membrane,Nucleus,Perikaryon,centriole,centrosome,cytoskeleton,growth cone,microtubule organizing center,perinuclear region,spindle
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 300-389 of human SIRT2 (NP_036369.2).
序列	IMGLGGGMDFDSKKAYRDVAWLGECDQGCLALAE LLGWKKELEDLVRREHASIDAQSGAGV PNPSTSASP KKSPPPAK DEARTTEREKPQ

靶点信息

研究背景	This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Several transcript variants are resulted from alternative splicing of this gene.
基因ID	22933
基因名	SIRT2
Swiss	Q8IXJ6
别名	SIRT2;SIR2;SIR2L;SIR2L2;sirtuin 2

产品验证



Western blot analysis of SIRT2 expressed in Rat brain using SIRT2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn