

IL17C Rabbit pAb

货号: AYP12650

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus IF: Mus musculus ChIP: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	21kDa
实测分子量	25kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,HT-29,Mouse small intestine,Rat lung
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 19-197 of human IL17C (NP_037410.1).
序列	HHDPSLRGHPHSHGTPHCYSAEELPLGQAPPHLLARGAKWQALPVALVSSLEAASHRGRHERPSATTQCPVLRPEEVLE ADTHQRSISPWRYRVDTDDEDRYPQKLAFEAELCRGCIDARTGRETAALNSVRLQLSLLVLRRRRCSRDRGSGGLPTPGAF TEFIHVPVGCTCVLPRSV

靶点信息

研究背景	The protein encoded by this gene is a T cell-derived cytokine that shares the sequence similarity with IL17. This cytokine was reported to stimulate the release of tumor necrosis factor alpha and interleukin 1 beta from a monocytic cell line. The expression of this cytokine was found to be restricted to activated T cells.
基因ID	27189
基因名	IL17C
Swiss	Q9P0M4
别名	IL17C;CX2;IL-17C

产品验证



Western blot analysis of IL17C expressed in Mouse small intestine, Rat lung, HeLa using IL17C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn