

— ABLYBIO, Help Your Research



# IL17C Rabbit pAb

货号: **AYP12650**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse,Rat  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  | <b>WB:</b> Mus musculus<br><b>IF:</b> Mus musculus<br><b>ChIP:</b> Mus musculus                      |
| 应用    | WB IF/ICC  |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000<br><b>IF/ICC:</b> 1:50 - 1:200   |
| 理论分子量 | 21kDa  |
| 实测分子量 | 25kDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | HeLa,HepG2,HT-29,Mouse small intestine,Rat lung  |
| 细胞定位  | Secreted   |
| 纯化    | Affinity purification  |


## 抗原信息

|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 19-197 of human IL17C (NP_037410.1). |
|------|--|

## 靶点信息

|       |   |
|-------|---|
| 研究背景  | The protein encoded by this gene is a T cell-derived cytokine that shares the sequence similarity with IL17. This cytokine was reported to stimulate the release of tumor necrosis factor alpha and interleukin 1 beta from a monocytic cell line. The expression of this cytokine was found to be restricted to activated T cells. |
| 基因ID  | 27189   |
| 基因名   | IL17C   |
| Swiss | Q9P0M4 ( <a href="https://www.uniprot.org/uniprotkb/Q9P0M4/entry">https://www.uniprot.org/uniprotkb/Q9P0M4/entry</a> )  |
| 别名    | IL17C,CX2,IL-17C,IL17C Rabbit pAb,Cytokine CX2  |

## 产品验证



Western blot analysis of IL17C expressed in Mouse small intestine, Rat lung, HeLa using IL17C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)