

— ABLYBIO, Help Your Research



BMP7 Rabbit pAb

货号: **AYP12677**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Mus musculus , Homo sapiens |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 49kDa |
| 实测分子量 | 49KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | SW480,293T,MCF7 |
| 细胞定位 | Secreted |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 70-251 of BMP7 (NP_001710.1). |
|------|---|

靶点信息

| | |
|-------|---|
| 研究背景 | This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer, which plays a role in bone, kidney and brown adipose tissue development. Additionally, this protein induces ectopic bone formation and may promote fracture healing in human patients. |
| 基因ID | 655 |
| 基因名 | BMP7 |
| Swiss | P18075 (https://www.uniprot.org/uniprotkb/P18075/entry) |
| 别名 | BMP7,OP-1,BMP7 Rabbit pAb,Osteogenic protein 1,OP1 |

产品验证



Western blot analysis of BMP7 expressed in SW480,293T,MCF7 using BMP7 Rabbit pAb at 1:1000. Second ary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)