

Phospho-POLR2A CTD-S2 Rabbit pAb

货号: **AYP12697**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB IP
推荐浓度	WB: 1:1000 - 1:2000 IP: 1:50 - 1:100
理论分子量	63kDa/217kDa
实测分子量	260kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A phospho specific peptide corresponding to residues surrounding S2 of human POLR2A CTD repeat YSPT SPS.
序列	SYSPT

靶点信息

研究背景	This gene encodes the largest subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. The product of this gene contains a carboxy terminal domain composed of heptapeptide repeats that are essential for polymerase activity. These repeats contain serine and threonine residues that are phosphorylated in actively transcribing RNA polymerase. In addition, this subunit, in combination with several other polymerase subunits, forms the DNA binding domain of the polymerase, a groove in which the DNA template is transcribed into RNA.
------	--

基因ID	5430
基因名	POLR2A
Swiss	P24928
别名	POLR2A;POLR2;POLRA;RPB1;RPBh1;RPO2;RPOL2;RpIILS;hRPB220;hsRPB1;Pol II

产品验证



Western blot analysis of Phospho-POLR2A CTD-S2 expressed in 293T using Phospho-POLR2A CTD-S2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn