

VDAC2 Rabbit pAb

货号: **AYP12708**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Danio rerio , Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	30kDa/31kDa/33kDa
实测分子量	31kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Rat brain
细胞定位	Mitochondrion outer membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-309 of human VDAC2 (NP_001171712.1).
序列	MSWCNELRLPALKQHSIGRGLESHITMCIPPSYADLGKAARDIFNKGFGFGLVKLDVKTSCSGVEFSTSGSSNTDTGKVT GTLETKYKWCEYGLTFTEKWNTDNTLGTEIAIEDQICQGLKLTFTDFTFSPNTGKKSGKIKSSYKRECINLGCDVDFDFAGPA IHGSAVFGYEGWLAGYQMTFDSAKSKLTRNNFAVGYRTGDFQLHTNVNDGTEFGGSIYQKVCEDLDTSVNLAWTSGTN CTRFGIAAKYQLDPTASISAKVNNSSLIGVGYTQTLRPGVKLTLSALVDGKSINAGGHKVGLELEA

靶点信息

研究背景	This gene encodes a member of the voltage-dependent anion channel pore-forming family of proteins that are considered the main pathway for metabolite diffusion across the mitochondrial outer membrane. The encoded protein is also thought to be involved in the mitochondrial apoptotic pathway via regulation of BCL2-antagonist/killer 1 protein activity. Pseudogenes have been identified on chromosomes 1, 2, 12 and 21, and alternative splicing results in multiple transcript variants.
------	--

基因ID	7417
基因名	VDAC2
Swiss	P45880
别名	VDAC2;POR

产品验证



Western blot analysis of VDAC2 expressed in Mouse brain,Rat brain using VDAC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn