

# TNFRSF10A Rabbit pAb

货号: **AYP12900**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse,Rat  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  | <b>WB:</b> Human fibrosarcoma cells , Homo sapiens<br><b>IHC:</b> Human colon cancer cell              |
| 应用    | <a href="#">WB</a> <a href="#">IF/ICC</a>  |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000<br><b>IF/ICC:</b> 1:50 - 1:100   |
| 理论分子量 | 50kDa  |
| 实测分子量 | 40-55kDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | K-562,SW480,MCF7,HeLa,DU145,Mouse lung,Rat lung  |
| 细胞定位  | Membrane,Single-pass type I membrane protein   |
| 纯化    | Affinity purification  |

## 抗原信息


|      |   |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 269-468 of human TNFRSF10A (NP_003835.3).   |
| 序列   | GGDPKCMDRVCFWRLGLLRGPGAEDNAHNEILSNADSLSTFVSEQMESQEPADLTGVTVQSPGEAQCLLGPAEAEQSGRRRLLVPANGADPTETLMLFFDKFANIVPFDSWDQLMRQLDLTKNEIDVVRAGTAGPGDALYAMLKWWNKTRNASIHTLLDALERMEERHAREKIQDLLVDSGKFIYLEDGTGSAVSLE |

## 靶点信息

|      |   |
|------|---|
| 研究背景 | The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. |
|------|---|

|       |   |
|-------|---|
| 基因ID  | 8797                                      |
| 基因名   | TNFRSF10A                                 |
| Swiss | O00220                                    |
| 别名    | TNFRSF10A;APO2;CD261;DR4;TRAILR-1;TRAILR1 |

## 产品验证



Western blot analysis of TNFRSF10A expressed in Mouse lung,Rat lung,K-562 using TNFRSF10A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30u g per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)