

— ABLYBIO, Help Your Research



HPD Rabbit pAb

货号: **AYP12932**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: HepG2 , Mus musculus
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	40kDa/44kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,Mouse liver,Mouse kidney,Rat liver,Rat kidney
细胞定位	cytosol,extracellular exosome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-393 of human HPD (N P_002141.1).
------	--

靶点信息

研究背景	The protein encoded by this gene is an enzyme in the catabolic pathway of tyrosine. The encoded protein catalyzes the conversion of 4-hydroxyphenylpyruvate to homogentisate. Defects in this gene are a cause of tyrosinemia type 3 (TYRO3) and hawkinsinuria (HAWK). Two transcript variants encoding different isoforms have been found for this gene.
基因ID	3242
基因名	HPD
Swiss	P32754 (https://www.uniprot.org/uniprotkb/P32754/entry)
别名	HPD,4-HPPD,4HPPD,GLOD3,HPPDASE,PPD,HPD Rabbit pAb,4-hydroxyphenylpyruvic acid oxidase

产品验证



Western blot analysis of HPD expressed in Mouse liver,Rat liver,HL-60 using HPD Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)