

NCR1 Rabbit pAb

货号: **AYP12933**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Mouse lung IF: Homo sapiens, Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	21kDa/22kDa/23kDa/32kDa/34kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HL-60
细胞定位	Cell membrane, Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-258 of human NCR1 (NP_001138929.2).
序列	SAQQQLPKPFIWAEPHFMVPKEKQVTICCGNYGAVEYQLHFEGSLFAVDRPKPPERINKVQFYIPDMNSRMAGQYSCIYRVGELWSEPSNLLDLVVTEMYDTPLSVHPGPEVISGEKVTFYCRDLDTATSMFLLKKEGRSSHVQRGYGKVAEFPLGPVTTAHRGTYRCFGSYNNHAWSPFPSEPVKLLVTGDIENTSLAPEDPTFPDTPWGTYLLTTETGLQKDHALWDHTAQNLLRM

靶点信息

研究背景	NKp46, along with NKp30 and NKp44, are activating receptors that have been collectively termed the natural cytotoxicity receptors (NCR). These receptors lack significant sequence homology to one another. They are expressed almost exclusively by NK cells and play a major role in triggering some of the key lytic activities of NK cells.
------	---

基因ID	9437
基因名	NCR1
Swiss	O76036
别名	CD335;LY94;NK-p46;NKP46;NCR1

产品验证



Western blot analysis of NCR1 expressed in HL-60 using NCR1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn