

RNF168 Rabbit pAb

货号: **AYP12941**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	IP: 293T , Homo sapiens WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:1000 - 1:2000
理论分子量	65kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,MCF7,Mouse brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 300-571 of human RNF168 (NP_689830.2).
序列	WLCACGAEWYHEGNVKTTPSNHGKELCVLSHERPKTRVPYSKETAVMPCGRTESEGCAPTSGVTQTNGNNTGETENEESCLLISKEISKRNQESSFEAVKDPFCFSARRRNVSPSSPDQEETEINFQKLIDLEHLLFERHKQEEQDRLLALQLQKEVDKEQMVPNRQKQSPDEYHLRATSSPPDKVLNGQRKNPKDGNFKRQTHTKHPTPERGSRDKNRQVSLKMQLKQSVNRRKMPNSTRDHCKVSKSAHSLQPSISQKSVFQMFQRCTK

靶点信息

研究背景	This gene encodes an E3 ubiquitin ligase protein that contains a RING finger, a motif present in a variety of functionally distinct proteins and known to be involved in protein-DNA and protein-protein interactions. The protein is involved in DNA double-strand break (DSB) repair. Mutations in this gene result in Riddle syndrome.
------	---

基因ID	165918
基因名	RNF168
Swiss	Q8IYW5
别名	RNF168;hRNF168

产品验证



Western blot analysis of RNF168 expressed in Mouse brain, HeLa, Jurkat using RNF168 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn