

— ABLYBIO, Help Your Research



KIAA0101 Rabbit pAb

货号: **AYP13086**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	7kDa/11kDa
实测分子量	12kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,K-562
细胞定位	Cytoplasm,Nucleus,perinuclear region
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 39-111 of human KIAA0101 (NP_055551.1).
------	---

靶点信息

研究背景	PCNA-binding protein that acts as a regulator of DNA repair during DNA replication. Following DNA damage, the interaction with PCNA is disrupted, facilitating the interaction between monoubiquitinated PCNA and the translesion DNA synthesis DNA polymerase eta (POLH) at stalled replisomes, facilitating the bypass of replication-fork-blocking lesions. Also acts as a regulator of centrosome number.
基因ID	9768
基因名	PCLAF
Swiss	Q15004 (https://www.uniprot.org/uniprotkb/Q15004/entry)
别名	PCLAF,KIAA0101,L5,NS5ATP9,OEATC,OEATC-1,OEATC1,PAF,PAF15,p15(PAF),p15/PAF,p15PAF,KIAA0101 Rabbit pAb,Hepatitis C virus NS5A-transactivated protein 9,Overexpressed in anaplastic thyroid carcinoma 1,PCNA-associated factor of 15 kDa,PCNA-clamp-associated factor

产品验证



Western blot analysis of KIAA0101 expressed in HT-29, K-562 using KIAA0101 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)