

ETS1 Rabbit pAb

货号: **AYP13095**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens,Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:100 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	25kDa/30kDa/40kDa/50kDa/55kDa
实测分子量	50KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 342-441 of human ETS1 (NP_005229 .1).
序列	LELLTDKSCQSFSISWTGDGWEFKLSDPDEVARRWGKRKNPKMNYEKLSRGLRYYYDKNIIHKTAGKRYVYRFVCDLQSL LGYTPEELHAMLVDVKPDADE

靶点信息

研究背景	This gene encodes a member of the ETS family of transcription factors, which are defined by the presence of a conserved ETS DNA-binding domain that recognizes the core consensus DNA sequence GGAA/T in target genes. These proteins function either as transcriptional activators or repressors of numerous genes, and are involved in stem cell development, cell senescence and death, and tumorigenesis. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.
------	--

基因ID	2113
基因名	ETS1
Swiss	P14921
别名	ETS-1;EWSR2;c-ets-1;p54;ETS1

产品验证



Western blot analysis of ETS1 expressed in Jurkat using ETS1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn