

RNF112 Rabbit pAb

货号: **AYP13099**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens,Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	68kDa
实测分子量	68KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat kidney
细胞定位	Membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 200-500 of human RNF112 (NP_009079.2).
序列	EGGRPRGGEASLQGCRWGANGLARGIWMWHPFLGKEGKKVAVFLVDTGDAMSPELSRETRIKLCALTTMLSSYQILS TSQELKDTDLDYLEMVHVAEVMGKHYGMVPIQHLLVDRDSSHPNKAGQGHVGNIFQRLSGRYPKVQELLQGKRARCC LLPAPGRRRMNQGHASPGDTDDDFRHLGAYVSDVLSAAPQHAKSRCQGYWNEGRAVARGDRRLTGQQLAQEIKNL SGWMGRTGPGFTSPDEMAAQLHDLRKVEAAKREFEYVRQQDVATKRIFSRVLPDTRNLLSTQ

靶点信息

研究背景	This gene encodes a member of the RING finger protein family of transcription factors. The protein is primarily expressed in brain. The gene is located within the Smith-Magenis syndrome region on chromosome 17.
------	--

基因ID	7732
基因名	RNF112
Swiss	Q9ULX5
别名	RNF112;BFP;ZNF179

产品验证



Western blot analysis of RNF112 expressed in Rat kidney using RNF112 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn