

# HCK Rabbit pAb

货号: **AYP13109**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	57kDa/59kDa
实测分子量	55-60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse spleen,Mouse lung,Rat spleen
细胞定位	Cell junction,Cell membrane,Cell projection,Cytoplasm,Cytoplasmic vesicle,Golgi apparatus,Lipid-anchor,Lysosome,Membrane,Nucleus,caveola,cytoskeleton,cytosol,cytosol,focal adhesion,podosome membrane,secretory vesicle
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 324-505 of human HCK (NP_001165600.1).
序列	LKSDEGSKQPLPKLIDFSAQIAEGMAFIEQRNYIHRDLRAANILVSASLVCKIADFGLARVIEDNEYTAREGAKFPIKWTAPEA INFGSFTIKSDVWSFGILLMEIVTYGRIPYPGMSNPEVIRALERGYRMPRENCPEELYNIMMRCWKNRPEERPTFEYIQSVL DDFYTATESQYQQQP

## 靶点信息

研究背景	The protein encoded by this gene is a member of the Src family of tyrosine kinases. This protein is primarily hemopoietic, particularly in cells of the myeloid and B-lymphoid lineages. It may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils. Multiple isoforms with different subcellular distributions are produced due to both alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) codon.
基因ID	3055
基因名	HCK
Swiss	P08631
别名	HCK;JTK9;p59Hck;p61Hck

## 产品验证



Western blot analysis of HCK expressed in Mouse spleen, Rat spleen, Raji using HCK Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)