

HAS3 Rabbit pAb

货号: **AYP13128**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	31kDa/62kDa
实测分子量	63KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	PC-3,Rat lung
细胞定位	Membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 67-281 of human HAS3 (NP_619515.1).
序列	HRRMRRAGQALKLPSRRGSVALCIAAYQEDPDYLRKCLRSAQRISFPDLKVMVVDGNRQEDAYMLDIFHEVLGGTEQA GFFVWRSNFHEAGEGETEASLQEGMDRVRDVVRASTFSCIMQKWGGKREVMYAFKALGDSVDYIQVCDSDTVLDPA CTIEMLRVLEEDPQVGGVGGDVQPPGKMAVEDDQVQAAQVRATEAWSVHQRHVSREQ

靶点信息

研究背景	The protein encoded by this gene is involved in the synthesis of the unbranched glycosaminoglycan hyaluronan, or hyaluronic acid, which is a major constituent of the extracellular matrix. This gene is a member of the NODC/HAS gene family. Compared to the proteins encoded by other members of this gene family, this protein appears to be more of a regulator of hyaluronan synthesis. Alternative splicing results in multiple transcript variants.
基因ID	3038
基因名	HAS3
Swiss	O00219
别名	HAS3

产品验证



Western blot analysis of HAS3 expressed in PC-3, Rat lung using HAS3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn