

# ATP13A2 Rabbit pAb

货号: **AYP13132**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens
应用	<a href="#">WB</a> <a href="#">IF/ICC</a>
推荐浓度	<b>WB:</b> 1:100 - 1:500 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	125kDa/128kDa
实测分子量	130KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	Lysosome,Membrane,Multi-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 67-210 of human ATP13A2 (NP_001135446.1).
序列	WKPLWGVRLRLRPCNLHAETLVIEIRDKEDSSWQLFTVQVQTEAIGEGSLEPSPQSAEDGRSQAAVAVGAVPEGAWKDTAQLHKSEEAKRVLRYLFFQGQRYIWIETQQAFYQVSLLDHGRSCDDVHRSRHLGLSLQDQMVRKAI

## 靶点信息

研究背景	This gene encodes a member of the P5 subfamily of ATPases which transports inorganic cations as well as other substrates. Mutations in this gene are associated with Kufor-Rakeb syndrome (KRS), also referred to as Parkinson disease 9. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	23400

基因名	ATP13A2
Swiss	Q9NQ11
别名	ATP13A2;CLN12;HSA9947;KRPPD;PARK9;SPG78

## 产品验证



Western blot analysis of ATP13A2 expressed in Mouse brain using ATP13A2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)