

Phospho-MAP2K4-T261 Rabbit pAb

货号: **AYP13160**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	44kDa/45kDa
实测分子量	46KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C6
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic phosphorylated peptide around T261 of human MAP2K4 (NP_003001.1).
序列	AKTRD

靶点信息

研究背景	This gene encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this family act as an integration point for multiple biochemical signals and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKKs, MAPKKs, and MAPKs. This protein is phosphorylated at serine and threonine residues by MAPKKKs and subsequently phosphorylates downstream MAPK targets at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver organogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicing results in multiple transcript variants.
------	--

基因ID	6416
基因名	MAP2K4
Swiss	P45985
别名	MAP2K4;JNKK;JNKK1;MAPKK4;MEK4;MKK4;PRKMK4;SAPKK-1;SAPKK1;SEK1;SERK1;SKK1

产品验证



Western blot analysis of Phospho-MAP2K4-T261 expressed in C6 using Phospho-MAP2K4-T261 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30u g per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn