

MED17 Rabbit pAb

货号: **AYP13208**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	15kDa/72kDa
实测分子量	80KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,A-431,Mouse testis,Rat testis
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 250-460 of human MED 17 (NP_004259.3).
序列	DLEGSAYIKVSIQKQAPDIGDLGTVNLFKRPLPKSKPGSPHWQTKLEAAQNVLLCKEIFAQLSREAVQIKSQVPHIVVKNQII SQPFPSLQLSISLCHSSNDKKSQKFATEKQCPEDHLYVLEHNLHLLIREFHKQTLSSIMMPHPASAPFGHKRMRLSGPQAF DKNEINSLQSSEGLLEKIIKQAKHIFLRSRAAATIDSLASRIEDPQI

靶点信息

研究背景	The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors.
基因ID	9440
基因名	MED17
Swiss	Q9NVC6
别名	MED17;CRSP6;CRSP77;DRIP80;TRAP80;SRB4

产品验证



Western blot analysis of MED17 expressed in Mouse testis,Rat testis,HeLa using MED17 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn