

— ABLYBIO, Help Your Research



F2RL3 Rabbit pAb

货号: **AYP13227**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	41kDa
实测分子量	41kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,HT-29,Raji,B cells,Mouse spleen,Mouse lung,Mouse skeletal muscle,Rat spleen
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-150 of human F2RL3 (NP_003941.2).
------	--

靶点信息

研究背景	Coagulation factor II (thrombin) receptor-like 3 (F2RL3) is a member of the large family of 7-transmembrane-region receptors that couple to guanosine-nucleotide-binding proteins. F2RL3 is also a member of the protease-activated receptor family. F2RL3 is activated by proteolytic cleavage of its extracellular amino terminus. The new amino terminus functions as a tethered ligand and activates the receptor. F2RL3 is activated by thrombin and trypsin.
基因ID	9002
基因名	F2RL3
Swiss	Q96RI0 (https://www.uniprot.org/uniprotkb/Q96RI0/entry)
别名	F2RL3,PAR4,F2RL3 Rabbit pAb,Coagulation factor II receptor-like 3,Thrombin receptor-like 3

产品验证



Western blot analysis of F2RL3 expressed in Mouse spleen,Rat spleen,A-549 using F2RL3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)