

RNF166 Rabbit pAb

货号: **AYP13255**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: Mus musculus
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	14kDa/17kDa/26kDa
实测分子量	26kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,SH-SY5Y,K-562,MCF7
细胞定位	cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100 to the C-terminus of human RNF 166 (NP_849163.1).
序列	GCNKKVTLAKMRVHISSCLKVQEQMANCPKFVPVVPVTSQIPISNIPNRSTFACPYCGARNLDQQELVKHCVESHRSDPNR VVCPICSAMPWGDPSYKSANFLQHLLHRHKFSYDTFVDYSIDEEAAAFQAALALSLEN

靶点信息

研究背景	E3 ubiquitin-protein ligase that promotes the ubiquitination of different substrates. In turn, participates in different biological processes including interferon production or autophagy. Plays a role in the activation of RNA virus-induced interferon-beta production by promoting the ubiquitination of TRAF3 and TRAF6. Plays also a role in the early recruitment of autophagy adapters to bacteria. Mediates 'Lys-29' and 'Lys-33'-linked ubiquitination of SQSTM1 leading to xenophagic targeting of bacteria and inhibition of their replication.
------	--

基因ID	115992
基因名	RNF166
Swiss	Q96A37
别名	RNF166

产品验证



Western blot analysis of RNF166 expressed in SW480,SH-SY5Y,K-562 using RNF166 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn