

CCDC92 Rabbit pAb

货号: **AYP13256**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB, IF: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	35kDa/36kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	SW480, BT-474
细胞定位	Cytoplasm, centriole, centrosome, cytoskeleton, microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-331 of human CCDC92 (NP_079416.1).
序列	MTSPHFSSYDEGPLDVSM AATNLENQLHSAQKNLLFLQREHASTLKGLHSEIRRLQQHCTDLTYELTVKSSEQTGDGTSK SSELKKRCEELEAQLKVKENENAELLKELEQKNAMITVLENTIKEREKKYLEELKAKSHKLTLLSSELEQRASTIAYLTSQLHAA KKKLMSSSGTSDASPSGSPVLASYKPAPPKDKLPETPRRRMKKSLSAPLHPEFEEVYRFGAESRKLKLLREPVDAMPDPTPF LLARESAEVHLIKERPLVIPPIASDRSGEQHSPAREKPKAHVGVVAHRIHHATPPQAQPEVKTLAVDQVNGGKVVVKHSGT DRTV

靶点信息

研究背景	Interferon-stimulated protein that plays a role in innate immunity. Strongly inhibits ebolavirus transcription and replication. Forms a complex with viral RNA-bound nucleocapsid NP and thereby prevents the transport of NP to the cell surface.
------	--

基因ID	80212
基因名	CCDC92
Swiss	Q53HC0
别名	CCDC92

产品验证



Western blot analysis of CCDC92 expressed in SW480, BT-474 using CCDC92 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn