

— ABLYBIO, Help Your Research



ERCC4 Rabbit pAb

货号: **AYP13262**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	42kDa/104kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,HT-1080
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 600-700 of human ERCC4 (NP_005227.1).
------	--

靶点信息

研究背景	The protein encoded by this gene forms a complex with ERCC1 and is involved in the 5' incision made during nucleotide excision repair. This complex is a structure specific DNA repair endonuclease that interacts with EME1. Defects in this gene are a cause of xeroderma pigmentosum complementation group F (XP-F), or xeroderma pigmentosum VI (XP6).
基因ID	2072
基因名	ERCC4
Swiss	Q92889 (https://www.uniprot.org/uniprotkb/Q92889/entry)
别名	ERCC4,ERCC11,FANCQ,RAD1,XFEPS,XPF,ERCC4 Rabbit pAb,DNA excision repair protein ERCC-4,DNA repair protein complementing XP-F cells,Xeroderma pigmentosum group F-complementing protein

产品验证



Western blot analysis of ERCC4 expressed in U-251MG,HT-1080 using ERCC4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)