

CKAP4 Rabbit pAb

货号: **AYP13291**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100
理论分子量	66kDa
实测分子量	66kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,MCF7,NCI-H460,Mouse testis
细胞定位	Cell membrane,Cytoplasm,Endoplasmic reticulum membrane,Single-pass type II membrane protein,cytoskeleton,perinuclear region
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 130-400 of human CKAP 4 (NP_006816.2).
序列	VLEEVQVRRSHQDFSRQREELGQGLQGVEQKVQSLQATFGTFESILRSSQHKQDLTEKAVKQGESEVSRISEVLQKLNKILKDLSDGIHVVKDARERDFTSLENTVEERLTELTKSINDNIAIFTEVQKRSQKEINDMKAKVASLEESEGNKQDLKALKEAVKEIQTSAKSREWDMEALRSTLQTMESDIYEVRELVSLKQEQQAFKEAADTERLALQALTEKLLRSEESVSRLPPEIRRLLEELRQLKSDSHGPKEDGGFRHSEAFE

靶点信息

研究背景	Mediates the anchoring of the endoplasmic reticulum to microtubules.
基因ID	10970

基因名	CKAP4
Swiss	Q07065
别名	CKAP4;CLIMP-63;ERGIC-63;p63

产品验证



Western blot analysis of CKAP4 expressed in Mouse testis,293T,HepG2 using CKAP4 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn