

— ABLYBIO, Help Your Research



IFNGR2 Rabbit pAb

货号: **AYP13305**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	37kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,F9
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200 to the C-terminus of human IFN GR2 (NP_005525.2).
------	--

靶点信息

研究背景	This gene (IFNGR2) encodes the non-ligand-binding beta chain of the gamma interferon receptor. Human interferon-gamma receptor is a heterodimer of IFNGR1 and IFNGR2. Defects in IFNGR2 are a cause of Mendelian susceptibility to mycobacterial disease (MSMD), also known as familial disseminated atypical mycobacterial infection. MSMD is a genetically heterogeneous disease with autosomal recessive, autosomal dominant or X-linked inheritance.
基因ID	3460
基因名	IFNGR2
Swiss	P38484 (https://www.uniprot.org/uniprotkb/P38484/entry)
别名	IFNGR2,AF-1,IFGR2,IFNGT1,IMD28,IFNGR2 Rabbit pAb,Interferon gamma receptor accessory factor 1,Interferon gamma receptor beta-chain,Interferon gamma transducer 1

产品验证



Western blot analysis of IFNGR2 expressed in HeLa, F9 using IFNGR2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)