

WBSCR22 Rabbit pAb

货号: **AYP13331**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100
理论分子量	24kDa/31kDa/33kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NCI-H460,A-549
细胞定位	Cytoplasm,Nucleus,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-281 of human WBSCR 22 (NP_059998.2).
序列	MASRGRRPEHGGPELFDYDETEARKYVRNSRMIDIQTRMAGRALELLYLLENKPCYLLDIGCGTGLSGSYLSDEGHYWVGL DISPAMLDEAVDREIEGDLGLGDMGQGIPIFKPGTFDGCISISAVQWLCNANKKSENPAKRLYCFFASLFSVLVRGSRVAVLQ LYPENSEQLELITTQATKAGFSGGMVVDYPNSAKAKKFFYLCLFSGPSTFIPEGLSENQDEVEPRESVFTNERFPLRMSRRG MVRKSRWVLEKKERHRRQGREVRPDTQYTGRKRKPRF

靶点信息

研究背景	This gene encodes a protein containing a nuclear localization signal and an S-adenosyl-L-methionine binding motif typical of methyltransferases, suggesting that the encoded protein may act on DNA methylation. This gene is deleted in Williams syndrome, a multisystem developmental disorder caused by the deletion of contiguous genes at 7q11.23. Alternatively spliced transcript variants have been found.
------	--

基因ID	114049
基因名	BUD23
Swiss	O43709
别名	BUD23;HASJ4442;HUSY-3;MERM1;PP3381;WBMT;WBSCR22

产品验证



Western blot analysis of WBSCR22 expressed in NCI-H460,A-549 using WBSCR22 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn