

# ING5 Rabbit pAb

货号: **AYP13337**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	<a href="#">WB</a> <a href="#">IF/ICC</a>
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	25kDa/27kDa
实测分子量	30KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human ING5 (N P_115705.2).
序列	MATAMYLEHYLDSIENLPCELQRNFQLMRELDQRTEDKKAIEDILAAEYISTVKTLSPDQRVERLQKIQNAYSKCKEYSDDK VQLAMQTYEMVDKHIRRLDADLARFEADLKDKMEGSDFFESSGGRGLKKGRGQKEKRGSRGRGRRTSEEDTPKKKKHK GGSEFTDTILSVHPSDVLDPVDPNEPTYCLCHQVSYGEMIGCDNPDCPIEFHFACVDLT

## 靶点信息

研究背景	This gene encodes a tumor suppressor protein that inhibits cell growth and induces apoptosis. This protein contains a PHD-type zinc finger. It interacts with tumor suppressor p53 and p300, a component of the histone acetyl transferase complex, suggesting a role in transcriptional regulation. Alternative splicing and the use of multiple promoters and 3' ends results in multiple transcript variants.
------	--

基因ID	84289
基因名	ING5
Swiss	Q8WYH8
别名	ING5;p28ING5

## 产品验证



Western blot analysis of ING5 expressed in 293T using ING5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)