

# TOX Rabbit pAb

货号: **AYP13363**

## 产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>IHC:</b> Mus musculus
应用	<a href="#">WB</a> <a href="#">IF/ICC</a> <a href="#">IHC</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200 <b>IHC:</b> 1:50 - 1:200
理论分子量	57kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 297-526 of human TOX (NP_055544.1).
序列	SMWDGLGEEQKQVYKKKTEAAKKEYLKQLAAYRASLVSKSYSEPVVDVKTSQPPQLINSKPSVFHGPSQAHSALYLSSHYY QQPGMNPHLTAMHPSLPRNIAPKPNQMPVTVSIANMAVSPPPPLQISPLHQHLNMQQHQPLTMQQPLGNQLPMQVQ SALHSPTMQQGFTLQPDYQTIINPTSTAAQVVTQAMEYVRSVSGCRNPPPQVVDWNNDYCSSGGMQRDKALYLT

## 靶点信息

研究背景	The protein encoded by this gene contains a HMG box DNA binding domain. HMG boxes are found in many eukaryotic proteins involved in chromatin assembly, transcription and replication. This protein may function to regulate T-cell development.
------	--

基因ID	9760
基因名	TOX
Swiss	O94900
别名	TOX;TOX1

## 产品验证



Western blot analysis of TOX expressed in CACO2, rat liver using TOX Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)