

EFTUD2 Rabbit pAb

货号: **AYP13366**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:100 IP: 1:20 - 1:50
理论分子量	105kDa/108kDa/109kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HepG2,HeLa,Mouse testis,Mouse brain,Mouse spleen,Mouse thymus,Mouse lung
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-205 of human EFTUD2 (NP_004238.3).
序列	MDTDLYDEFGNYIGPELDSDEDDDELGRETKLDDEMDDDDDDDDVGDHDDHDPGMEVVLHEDKKYYPTAEEVYGPEV ETIVQEEDTQPLTEPIIKPVKTKKFTLMEQTLPTVYEMDFLADLMDNSELIRNVTLCGHLHHGKTCFVDCLIEQTHPEIRKRY DQDLCYTDILFTEQERGVGKSTPVTVVLPDTKGKSYLFNIMDT

靶点信息

研究背景	This gene encodes a GTPase which is a component of the spliceosome complex which processes precursor mRNAs to produce mature mRNAs. Mutations in this gene are associated with mandibulofacial dysostosis with microcephaly. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	9343
基因名	EFTUD2
Swiss	Q15029
别名	EFTUD2;MFDGA;MFDM;SNRNP116;Snrp116;Snu114;U5-116KD

产品验证



Western blot analysis of EFTUD2 expressed in Mouse testis, Jurkat, HepG2 using EFTUD2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn