

RARRES2 Rabbit pAb

货号: AYP13377

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	18kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,BT-474,HeLa,Mouse liver
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 17-163 of human RARRES2 (NP_002880.1).
序列	VGVAELTEAQRRLQVALEEFHKHPPVQWAFQETSVESAVDTPFPAGIFVRLFEKLLQQTSCRKRDWKKPECKVRPNGRKR KCLACIKLGSSEDKVLGRLVHCPIETQVLREAEHQETQCLRVQRAGEDPHSFYFPGQFAFSKALPRS

靶点信息

研究背景	This gene encodes a secreted chemotactic protein that initiates chemotaxis via the ChemR23 G protein-coupled seven-transmembrane domain ligand. Expression of this gene is upregulated by the synthetic retinoid tazarotene and occurs in a wide variety of tissues. The active protein has several roles, including that as an adipokine and as an antimicrobial protein with activity against bacteria and fungi.
------	---

基因ID	5919
基因名	RARRES2
Swiss	Q99969
别名	RARRES2;HP10433;TIG2

产品验证



Western blot analysis of RARRES2 expressed in Mouse liver,293T,BT-474 using RARRES2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn