

NNMT Rabbit pAb

货号: **AYP13402**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	29kDa
实测分子量	28KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-264 of human NNMT (NP_006160.1).
序列	MESGFTSKDTYLSHFNPRDYLEKYYKFGSRHSAESQILKHLKLNLFKIFCLDGVKGDLLIDIGSGPTIYQLLSACESFKEIVT DYSDQNLQELEKWLKKEPEAFDWSPVVTYVCDLEGNRVKGPEKEEKLQAVKQVLKCDVTQSQPLGAVPLPPADCVLST LCLDAACPDLPYCRALRNLSLLKPGGFLVIMDALKSSYIMIGEQQFSSPLGREAVEAAVKEAGYIEWFEVISQSYSST MANNEGLFSLVARKLSRPL

靶点信息

研究背景	N-methylation is one method by which drug and other xenobiotic compounds are metabolized by the liver . This gene encodes the protein responsible for this enzymatic activity which uses S-adenosyl methionine as the methyl donor.
------	---

基因ID	4837
基因名	NNMT
Swiss	P40261
别名	NNMT

产品验证



Western blot analysis of NNMT expressed in A-549 using NNMT Rabbit pAb at 1:1000. Secondary antibody : HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn